



THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION. CONTAINS 2,4-D, MECOPROP-p AND DICAMBA
Weeds contacted by this product gradually die and disappear from the lawn.

ACTIVE INGREDIENTS:
Isocetyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid* ... 1.077%
2,4-Dichlorophenoxyacetic Acid* 0.146%
(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**‡ 0.287%
Dicamba (3,6-Dichloro-o-Anisic Acid)*** 0.070%
OTHER INGREDIENTS: 98.420%
TOTAL: 100.000%

Isomer Specific Method, Equivalent to:
*2,4-Dichlorophenoxyacetic Acid (Total) 0.860%
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid 0.287%
***3,6-Dichloro-o-Anisic Acid 0.070%
‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p

EPA REG. NO.228-413-57131
EPA EST. NO.57131-WI-1

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are any waterproof materials. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators, flaggers, and other handlers must wear:

• long-sleeved shirt and long pants, • shoes and socks, plus • chemical-resistant gloves (except for applicators using groundboom equipment)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS
This pesticide may be toxic to fish and aquatic invertebrates and may be adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not enter or allow people (or pets) to enter the treated area until dusts have settled. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.

This product is a stable mixture of four herbicides. It is for use on Ornamental Turf lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and controls many broadleaf weeds such as:

Alder	Common chickweed	Horsenettle	Prostrate pigweed	Tarweed
Annual yellow	Common mullein	Horseail	Prostrate spurge	Thistle
sweet clover	Common sowthistle	Indiana mallow	Prostrate vervain	Tick trefoil
Artichoke	Corn	Ironweed	Puncture vine	Toadflax
Aster	Chamomile	Jewelweed	Purslane	Trailing Crownvetch
Austrian fieldress	Creeping jenny	Jimsonweed	Ragweed	Tumble mustard
Bedstraw	Crimson clover	Kochia	Rumex	Tumble pigweed
Beggartick	Croton	Knawel	Redroot pigweed	Tumbleweed
Biden	Cudweed	Knotweed	Red sorrel	Velvet leaf
Bindweed	Curly dock	Lambsquarter	Redstem filaree	Vernice mallow
Bird vetch	Curly indigo	Lespedeza	Rough cinquefoil	Veronica
Bitterweed	Dandelion	Locoweed	Rough fleabane	Vervain
Bitter wintercress	Dead nettle	Lupine	Roundleafed marigold	Vetch
Black-eyed Susan	Dock	Mallow	Rush	Virginia buttonweed
Black medic	Dogbane	Marshelder	Russian pigweed	Virginia creeper
Black mustard	Dogfennel	Matchweed	Russian thistle	Virginia pepperweed
Blackseed plantain	Elderberry	Mexicanweed	St. Johns wort	Wayleaf bullthistle
Blessed thistle	English daisy	Milk vetch	Scarlet pimpernel	Western dematis
Blue lettuce	Fall dandelion	Milkweed bloodflower	Scotch thistle	Western salsify
Blue vervain	False dandelion	Mouseear chickweed	Sheep sorrel	White clover
Box elder	False flax	Morningglory	Shepherdspurse	White mustard
Bracted plantain	False sunflower	Muscleweed	Slender plantain	Wild mustard
Brassbuttons	Fiddleneck	Musk thistle	Smallflower galinsoga	Wild aster
Bristly oxtongue	Field bindweed	Mustard	Smoothweed	Wild buckwheat
Broadleaf dock	Field pansy	Narrowleaf plantain	Smooth dock	Wild carrot
Broadleaf plantain	Flea bane (daisy)	Narrowleaf vetch	Smooth pigweed	Wild four-o'clock
Broomweed	Flaxweed	Orange hawkweed	Nettle	Wild garlic
Buckhorn	Florida betony	Oxalis	Southern wild rose	Wild geranium
Buckhorn plantain	Florida pusley	Oxeye daisy	Sowthistle	Wild lettuce
Bulbous buttercup	Frenchweed	Parsley-piert	Spanishneedle	Wild marigold
Bull nettle	Galinsoga	Parsnip	Spatterdock	Wild onion
Bull thistle	Garlic mustard	Pearwort	Speedwell	Wild parsnip
Burdock	Goathead	Pennywort	Spiny Amaranth	Wild radish
Burning nettle	Goatsbeard	Pennywort	Spiny cocklebur	Wild rape
Bur ragweed	Goldenrod	Peppergrass	Spotted catsear	Wild strawberry
Burweed	Ground ivy	Pepperweed	Spotted knapweed	Wild sweet potato
Buttercup	Gumweed	Pigweed	Spurred spurge	Wild vetch
Canada thistle	Hairy bittercress	Pineywoods bedstraw	Spurge	Willow
Carolina geranium	Hairy fleabane	Plains coreopsis	Spurweed	Witchweed
Carpetweed	Hawkweed	Plantain	Stinging nettle	Woolly morningglory
Catchweed bedstraw	Healall	Poison hemlock	Stinkweed	Woodsorrel
Catsear	Hearthleaf drymar	Poison ivy	Stitchwort	Woolly croton
Catnip	Hedge bindweed	Poison oak	Strawberry clover	Woolly plantain
Chickweed	Hedge mustard	Pokeweed	Sumac	Wormseed
Chicory	Hemp	Poorjoe	Sunflower	Yarrow
Cinquefoil	Henbit	Povertyweed	Sweet clover	Yellow rocket
Clover	Hoary cress	Prickly lettuce	Tail nettle	Yellow flower
Cockle	Hoary plantain	Prickly sida	Tansy mustard	pepperweed
Cocklebur	Hoary vervain	Primrose	Tansy ragwort	
Coffeebean	Hop clover	Prostrate knotweed		
Coffeeweed				

WHEN TO APPLY: Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 50 pound bag should be used on 12,500 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound per 250 square feet.

CALIBRATION INSTRUCTIONS:

To provide proper distribution, calibrate spreader before application. For 4.0 pounds of product per 1,000 square feet of turfgrass. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not stack more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

This product is protected by U.S. Patent No. 6,579,831. No license granted under any non-U.S. Patent(s). Viper is a Registered Trademark of Nufarm Americas Inc.

(RV20170614)

Net Wt. 50 lbs. (22.7 kg)

Distributed and Guaranteed by Eau Claire Co-op Oil Co.
P.O. Box 837, Eau Claire, WI 54702
www.ecgrow.com