


MILLENNIUM ULTRATM WEED & FEED



ACTIVE INCOEDIENTS

ACTIVE INGREDIENTS:		
Clopyralid Acid**		
TOTAL: BY ISOMER SPECIFIC METHOD, EQUIVALENT TO:		
	0.071%	
EPA REG. NO. 228-343-57131		
EPA EST. NO. 57131-WI-1	KEEP OUT OF REACH OF CHILDREN	
IN NEW YORK STATE, THIS PRODUCT MAY ONLY BE		
SOLD TO AND USED BY PROFESSIONAL APPLICATORS	CAUTION	
AND IS PROHIBITED FROM SALE AND USE IN NASSAU		
AND SUFFOLK COUNTIES. IN CALIFORNIA, OREGON,	SEE BACK PANEL FOR FIRST AID AND	
AND WASHINGTON, TURFGRASS AND LAWN USES ARE RESTRICTED TO GOLF COURSES ONLY. This product	ADDITIONAL PRECAUTIONARY STATEMENTS	
cannot be used on Agricultural, Forest sites or Right-of-way		
carnot be used on Agricultural, i orest sites of hight-or-way	For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300 For Medical Emergencies Only, call (877) 325-1840	
	For weuldar Entergencies Only, Call (677) 323-1040	
	UTIONARY STATEMENTS UMANS AND DOMESTIC ANIMALS CAUTION	
Causes moderate eye irritation. Avoid contact with eye: PERSONAL PROTECTIVE EQUIPMENT (PPE): All loaders, applicators, and other handlers must w		
	edi.	
 long-sleeved shirt and long pants 		
shoes plus socks		
chemical-resistant gloves made of any waterproof material		
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot		
water. Keep and wash PPE separately from other laun	dry.	
USER SA	FETY RECOMMENDATIONS	
Users Should:		
• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.		
 Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on 		
skin, wash immediately with soap and water.		
	Wash the outside of gloves before removing. As soon as possible, wash	
thoroughly and change into clean clothing.	. Wash are balance of gives before removing. As soon as possible, Wash	
u lo lo ugi liy and change into ciedri cioti ling.		

FIRSTAID	
IF IN EYES:	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with it's labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Not for use on turf grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is for use on Ornamental Turf Lawns (Industrial and Institutional), Parks Cemeteries, Athletic and Recreational Sports Fields, Golf Courses (Fairways, Aprons, Tees* and Roughs) and similar turf areas. *Excluding Bentgrass Tees.

The maximum number of broadcast applications per treatment site is 2 per year

Artichoke

Bedstraw

Biden

Black

Beggartick

Bindweed

Bird vetch

Ritterweer

Black Mustard

Blessed th

Blue lettuce

Brassbuttons

Broadleaf dock

Broomweed

Buckhorn

Bull nettle

Bull thistle

Burning nettle

Bur raqweed

Canada thistle

Carpetweed

Catsear

Catnip

Chicorv

Clover

Cockle

Cinquefoil

Cocklebur

Coffeebear

Coffeeweer

Chickweed

Burdock

Burweed

Buttercup

ervair

Aster

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

Do not use on Centipede, Dichondra, and St. Augustine. Do not use this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants. Do not apply when air temperature exceeds 90°F. Avoid overlapping.

Do not use on residential turf. Do not collect grass clippings for mulch or compost. Do not send clippings to a compost facility. Applicator must give notice to landowners/property managers to not use grass clippings for mutching or composting. WEEDS CONTROLLED

This product is designed to promote lush green growth in lawins and control many broadleaf weeds such as: Alde

Common chickweed Horseta Annual yellow sweet clover Common mullein ndiana n Common sowthistle Tronweed Corn Chamomi .lewelweer Austrian fieldcress Creeping i .limsonweei nawel Knotweed Lambsquarte Lespedeza upine Mallow Marshelde Matchweed Mexicanweed Black-seed plantai Elderbe English daisy Milk vetch Fall dandelion Milkweed bloodflower False dandelior Muawort False flax Morningglory False sunflower cted plantair Mousear chickweed Fiddleneck Musk thistle Bristly oxtongue Field bindweed Mustard Field pansy Narrowleaf plantain Narrowleaf vetch Broadleaf plantain Flea bane (daisy) Flixweed Nettle Florida betony Orange hawkweed Buckhorn plantair Florida pusley Oxalis Bulbous buttercup Frenchweed Oxeye daisy Galinsoga Parsley-piert Garlic mustard Parsnip Goathead Pearlwort Goatsheard Pennycress Goldenrod Pennywort Ground ivv Peppergrass Gumweed Pepperweed Hairv bittercress Piaweed Hairy fleabane Pineywoods bedstraw Carolina geranium Hawkweed Plains coreopsis Catchweed Bedstraw Healall Plantain Heartleaf drymary Poison hemlock Hedge bindweed Poison ivv Hedge mustard Poison oak Hemp, Henbit Pokeweed Hoary cress Poorjoe Hoary plantain Povertyweed Prickly lettuce Hoarv vervain Prickly sida Honevsuckle Hop clover Primrose Prostrate knotweer Horsenettle

Prostrate pigweed had rostrate spurge Tarweed trate vervair Thistle Tick trefoil Puncture vine Toadflax Trailing Crownyetch Ranweed Red clover Tumble mustard Redroot pigweed Tumble pigweed Red sorrel Tumbleweed Redstem filaree Velvet leaf Rough cinquefoil Venice mallow Rough fleabane Veronica Roundleafed marigold Vervain Rush Vetch Virginia buttonweed Russian pigweed Russian thistle Virginia creeper St. Johnswort Virginia pepperweed Scarlet pimpernel Wavvleaf bullthistle Scotch thistle Western clematis Sheep sorrel Western salsify White clover Shepherdspurse Slender plantair White mustard Smallflower galinsoga Wild aster Smartweed Wild buckwheat Wild carrot Wild four-o'-clock Wild garlic Southern wild rose Wild geranium Wild lettuce Wild marigold Wild mustard Wild onion Wild parsnip Wild radish Wild rape Wild strawberry Spotted knapweed Wild sweet potato Wild vetch Willow Witchweed Woolly croton Stitchwort Woolly morning-glory Strawberry clover Woolly plantain Sumac Woodsorrel Sunflower Wormseed Sweet clover Yarrow Tall nettle Yellow rocket Tall vervain Yellowflower pepperweed

WHEN TO APPLY: Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to listed Warm Season Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 50 pound bag should be used on 12,500 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for a slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound per 250 square feet.

s also acceptable.			
SPREADER SETTINGS			
4 lbs Per 1,000 Sq. Ft.			
Lesco Rotary	F-G		
Spyker 24-22	4		
Scott's R8A	1		
Prizelawn CB Rotary	4		
Earthway 2400 Rotary	17		
Lely Broadcast	3.4		

Sweep (co not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas oil to reduce corrosion.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not stack more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides. PRODUCT DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER. EITHER EXPRESS OR IMPLIED. OR BY USAGE OF TRADE. STATUTORY OR OTHERWISE. WITH REGARD TO THE PRODUCT SOLD. INCLUDING. BUT NOT LIMITED TO MERCHANTABILITY. FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS. MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD. INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BYWAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER. IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY. CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF OR RISING OUT OF THE MISUSE. OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER. USER. OR ITS CUSTOMERS TO THE EXTENT CONSISTENTWITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Millennium Ultra is a Registered Trademark of Nufarm Americas Inc.

```
(RV012816N)
```

Distributed and Guaranteed by: Eau Claire Co-op Oil Co. P.O. Box 837, Eau Claire, WI 54702 www.ecgrow.com

NET WT. 50 LBS. (22.7 Kg)

Smooth dock Smooth pigweed Sneezeweed

Sowthistle Spanishneedle Spatterdock Speedwell Spiny amaranth Spiny cocklebu Spotted catsear Spotted spurge Spurge Spurweed Stinging nettle Stinkweed

Tansy mustard and other broadleaf Tansy ragwort weeds